



Brief Communication

Late Pleistocene foot infection in *Dama mesopotamica* from Tabun B (Mount Carmel, Israel)Ana B. Marín-Arroyo^{a,*}, Francisco Gil Cano^b, Mark Lewis^c^a Instituto Internacional de Investigaciones Prehistóricas de Cantabria, Universidad de Cantabria, Avda. de los Castros, s/n, 39005 Santander, Spain^b Departamento de Anatomía y Anatomía Patológica Comparadas, Facultad de Veterinaria, Universidad de Murcia, Avda. Teniente Flomesta, 5, 30003 Murcia, Spain^c Department of Earth Sciences, Natural History Museum, London SW7 5BD, United Kingdom

ARTICLE INFO

Article history:

Received 13 March 2013

Received in revised form 21 August 2014

Accepted 25 August 2014

Keywords:

Fallow deer

Phalanx

Osteitis

Footrot

Tabun

Late Pleistocene

ABSTRACT

Tabun is one of the most important Palaeolithic sites in the near East, with levels dating from the Lower through to the Upper Palaeolithic. The faunal collection from Tabun Cave (Israel) was recovered by Dorothy Garrod during archaeological excavations carried out during the 1920–30s in Mount Carmel. Since then this collection has been housed at The Natural History Museum, London. In this brief communication, a frequently occurring pathology in the phalanges of fallow deer (*Dama mesopotamica*) from Level B is reported. The pathology consists of bone resorption on the cortex of the axial surface of the proximal phalanges, most likely as a result of a localized osteitis produced by an infection, initially on the hoof, and later on the phalanges and even metapodials. These pathologies were probably caused by bacterial infection, possibly linked to environmental and climatic conditions at the site.

© 2014 Elsevier Inc. All rights reserved.

1. Introduction

Tabun contains evidence of human hunting from the late Lower Palaeolithic to modern times (Stiner et al., 2010), and is one of the most important Middle Palaeolithic sites in the near East (Shea, 2003). The faunal collection from Tabun Cave (Israel) was recovered by Dorothy Garrod during archaeological excavations carried out during the 1920–30s in Mount Carmel (Garrod and Bate, 1937). Dorothea Bate (1937) was first to identify the macro-mammal assemblage through the whole stratigraphic sequence. In the 1980s, Garrard (1981) reviewed the previous palaeontological work and, more recently, Marín-Arroyo (2013a, 2013b) undertook a taphonomic analysis of the Mousterian levels. Through the study of bone modifications, several identical lesions were observed in fallow deer remains, specifically in the proximal phalanges. The particular pathologies reported here are uncommon within the archaeological record but interesting enough to understand their cause among several artiodactyl species. Currently, this collection is curated at the Department of Earth Sciences at the Natural History Museum, London.

2. Results

Fallow deer is the most common taxon found in level B of Tabun Cave, with all anatomical elements represented within the deposit and dated to between 130 and 70 kyr (MIS 5). During the taphonomic study of the large mammals carried out by Marín-Arroyo (2013a, 2013b), some particularly deep cortical bone lesions were observed with the naked eye on the axial surfaces of several proximal phalanges and later were examined with a binocular microscope. It was notable through microscopy that these lesions were not caused anthropogenically, but were the result of a pathology. From a NISP of 97 elements, 15% of fused proximal phalanges showed these alterations ($n = 15$). In the phalanges, the individual bone lesion was always located on the cortex of the axial surface of the proximal phalanges, just distal to the fusion line of the proximal epiphysis (Fig. 1). The diameter of these lesions was, on average, 4–5 mm in width and 2–3 mm depth (Fig. 2). On one phalanx, new bone formation was identified affecting the complete bone (Fig. 3). Unfortunately, it was not possible to discriminate the sex of the animals and whether or not phalanges belonged to fore or hind feet.

The appearance of the cortical bone lesion on the axial surfaces of the proximal epiphyses could be a localized osteitis, probably as a result of contagious bacterial infection, which usually starts on the hoof. Currently, this infection, known as footrot, is

* Corresponding author. Tel.: +34 942202016.

E-mail address: marinab@unican.es (A.B. Marín-Arroyo).



Fig. 1. Proximal phalanges of fallow deer from Tabun B. The pathology is observed in the cortex of the axial surfaces of the proximal phalanges. Photo courtesy of the NHM, London.



Fig. 2. The image shows a close view of the osteitis on the proximal part of the axial surface of the phalanx. Scale shows millimeters. Photo courtesy of the NHM, London.

caused by the microbial activity of two bacteria, *Fusobacterium necrophorum* and *Dichelobacter nodusus* (Ginn et al., 2007). The former is regularly present in soil and manure, and the latter, when present, is the bacterium that causes the problem. Unfortunately, it has not been possible to prove whether or not these bacteria were those involved in the infection during MIS 5, the period when the cave deposit accumulated. Nevertheless, specific environmental conditions such as warmth and moisture are important factors in this disease; as the hooves soften they become susceptible to colonization by the bacteria responsible in the development of footrot. In the same way, rocky soils also increase the likelihood of infection, due to the risk of trauma to the hoof.

Although the infection grows within the hoof, it spreads via the soft tissues to the surface of the phalanges where the infection then spreads along the bone surface to the metapodials in severe cases. When the climatic and geological conditions described above exist, the activity of *F. necrophorum* proliferates affecting the interdigital dermis, causing soft tissue inflammation and osteoclast-mediated lesion of the surface of metapodials bones in severe cases (Fernández et al., 1996; Thompson, 2007).

In the case of Tabun B, the pattern of bone surface lesion is compatible with one caused by a localized bacterial periostitis and one that has also involved the insertion of the collateral ligament on the axial surface of the proximal phalanx. Avulsion of the insertion of this ligament would have induced the formation of the enthesophyte observed at this location in the proximal phalanx of these specimens.



Fig. 3. Proximal phalanx of fallow deer from Tabun B with new bone formation across the entire shaft. From left to right: medial, lateral, dorsal and plantar views. Photo courtesy of the NHM, London.

3. Discussion and conclusions

Currently, footrot is often found among species of the order Artiodactyla, such as cervids and bovids, when particular climatic and landscape conditions are present; it has also been reported in pigs (Ginn et al., 2007). During a large part of MIS 5, the Levant region had a warm climate with high humidity (Bar-Matthews et al., 2003; Marín-Arroyo, 2013a; Fig. 5), which at times would have allowed the hooves of fallow deer to soften, permitting microorganism activity with subsequent infection in the interdigital spaces, and later spreading to the bone. Additionally, the topographic relief of the grassland feeding locations and of the area surrounding Mount Carmel would have facilitated the occurrence of injuries within the hoof. Today, footrot is a devastating disease, initially affecting the hooves, with lameness as the first visible signal within the herds. When the infection increases, necroses appear in the soft tissues, causing lesions in the ligaments and later in the adjacent bones (phalanges and metapodials). These lesions exacerbate the lameness of the animal which reduces their survivorship and ability to search for food, making them more susceptible to predators or natural traps, as was the case with Tabun B (Marín-Arroyo, 2013a,b).

Acknowledgements

Thanks are due to Adrian M. Lister and Andy Currant (Natural History Museum, London) for making the Tabun collection available for study. The comments by Eleanor Clarke helped to improve this manuscript. Ana B. Marín-Arroyo has a contract of the Ramón y Cajal Research Program (RYC-2011-00695), at the Instituto Internacional de Investigaciones Prehistóricas de Cantabria, Universidad de Cantabria. This Research is supported with funding

from the Spanish Ministry of Economy and Competitiveness Ref. No.: HAR2012-33956 and European Union through FP7-PEOPLE-2012-CIG (Ref. No.: 322112).

References

- Bar-Matthews, M.A., Ayalon, A., Gilmour, M., Matthews, A., Hawkesworth, C.J., 2003. Soreq and Peqiin Caves, Israel Speleothem Stable Isotope Data, IGBP PAGES/World Data Center for Paleoclimatology Data Contribution Series #2003-061. NOAA/NGDC Paleoclimatology Program, Boulder, CO.
- Bate, D.M.A., 1937. Paleontology: the fossil fauna of Wady el-Mughara caves. In: Garrod, D.A.E., Bate, D.M.A. (Eds.), *The Stone Age of Mount Carmel, Part 2*. Clarendon Press, Oxford, pp. 137–240.
- Garrard, A., (PhD dissertation unpublished) 1981. *Man–animal–plant relationship during the Upper Pleistocene and early Holocene of the Levant*. University of Cambridge, Cambridge.
- Garrod, D.A.E., Bate, D.M.A., 1937. *The Stone Age of Mount Carmel. Excavations at the Wadi Mughara, vol. 1*. Clarendon Press, Oxford.
- Ginn, P.J., Mansell, K.L., Rakich, P.M., 2007. Skin and appendages. Fungal diseases of skin. In: Grant Maxie, M. (Ed.), *Jubb, Kennedy & Palmer's Pathology of Domestic Animals, vol. 1(5)*, fifth ed. Elsevier Saunders, pp. 694–709, <http://www.sciencedirect.com/science/book/9780702028236>
- Fernández, M., Manzanera, Y., Serrano, E., 1996. *Pedero del ovino y caprino*. *Mundo ganadero* 81, 46–50.
- Marín-Arroyo, A.B., 2013a. Palaeolithic human subsistence in Mount Carmel (Israel). A taphonomic assessment of Middle and Early Upper Palaeolithic faunal remains from Tabun, Skhul and el-Wad. *Int. J. Osteoarchaeol.* 23 (3), 254–273.
- Marín-Arroyo, A.B., 2013b. New opportunities for previously excavated sites: palaeo-economy as a human evolutionary indicator at Tabun Cave (Israel). In: Clark, J., Speth, J. (Eds.), *Zooarchaeology and Modern Human Origins, Human Hunting Behavior during the Later Pleistocene*. Vertebrate Paleobiology and Paleoanthropology Series. Springer, London, pp. 59–75.
- Shea, J.J., 2003. The Middle Paleolithic of the East Mediterranean Levant. *J. World Prehist.* 17 (4), 313–394.
- Stiner, M.C., Gopher, A., Barkai, R., 2010. Hearth-side socioeconomics, hunting and paleoecology during the late Lower Paleolithic at Qesem Cave, Israel. *J. Hum. Evol.* 60, 213–233.
- Thompson, K., 2007. Bone and joints. Inflammatory diseases of bones. In: Grant Maxie, M. (Ed.), *Jubb, Kennedy & Palmer's Pathology of Domestic Animals, vol. 1(1)*, fifth ed. Elsevier Saunders, pp. 92–105, <http://www.sciencedirect.com/science/book/9780702028236>